

PLEASE ATTACE	PATIENT DE	MOGRAPHIC ANL	INSUKAI	NCE INFORIVI	ATION							
PATIENT DEMOG	RAPHICS											
PATIENT												
NAME				DOB				in	WEIG	інт	kg	
Posses II						ALLERGIES						
DIAGNOSIS PHONE #						Include OTC/herbal						
PRIMARY						PRIMARY						
INSURANCE		INSURANCE #										
EMERGENCY												
CONTACT						PHONE	‡					
THERAPY INFORM	MATION											
ORDERING												
PROVIDER							#					
FOLLOWING												
PROVIDER							#					
EXISTING	IG □ Central Line (Tunneled/Non-tunneled) □ Peripheral IV					☐ Port	Needle size	e	Accessed			
IV ACCESS	☐ Midline:	lumen	□ Other:									
2201/1252 2225	-				_							
MEDICATION		Door	Bours	Facousies	Tuspany	Leneru	OHANETY	C=+D	- D	Crop Dar		
	DRUG	Dose C rea //er	ROUTE	FREQUENCY	THERAPY	LENGIH	QUANTITY	STAR	T DATE	STOP DAT	E	
		□ Cubicin® 6 mg/kg IV q 24 hours			#QS							
	□ Invanz® 1 gram		IV	q 24 hours			#QS					
	☐ Vancomyc	in 1000 mg	IV q 12 hours				#QS					
	□ Ceftriaxon	e 2 grams	IV	q 24 hours			#QS					
			IV	q			#QS					
				q			#QS					
		Flu	ush with 0.9		s) before a	and after		. followed	by hepar	in lock		
	□ Peripheral IV (PIV) Flush with 0.9% NaCl (5 mLs) before and after medication, followed by heparin lock (10 units/mL) 5 mLs as a final lock (SASH) # QS											
	☐ Midline, PICC, Central Flush with 0.9% NaCl (10 mLs) before and after medication, followed by heparin lock											
FLUSH	Venous Catheters (10 units/mL) 5 mLs after completion of medications (SASH); Flush additional lumen with											
PROTOCOL (Select one)	(Single, double, triple lumen) 0.9% NaCl (10 mLs) followed by heparin lock (10 units/mL) 5 mLs once daily #QS											
(Select offe)	Flush port with 0.9% NaCl (10 mLs) before and after medications, followed by heparin lock											
	(100 units/mL) 5 mLs after completions of medications #QS											
	□ Other:											
SUPPLIES	□ Supplies and pumps necessary to maintain and administer medication											
	☐ Anaphylax	ris Kit: Diphenhydrar	mine 50 mg	(1 vial); Epine	phrine 1:	1000 (2 v	ials); Suppli	es for adn	ninistratio	n		
ANAPHALYXIS	Allergic response - As per provider order: Diphenhydramine 50 mg slow IV push over 2-3 minutes											
KIT	_	axis - As per provide				-	•			M injection;		
	Epinephi	rine 1:1000 solution	: 0.4 mg (0.	.4 mL) subcuta	neous inje	ection; If	needed, ma	ıy repeat i	n 20 minu	tes times 1 do	ose	
	□ IV therapy administration by skilled nursing personnel											
	□ Patient education on administration of IV therapy performed during skilled nursing visit											
IV ACCESS	□ Peripheral IV site to remain on condition site viable; Restart upon any level of pain/tenderness, changes in skin color or											
MAINTENANCE	temperature, edema, induration, fluid leakage/drainage, or other abnormality and as needed to maintain therapy access											
	□ Subcutaneous port re-access every 7 days and as needed at home or clinic											
	□ Dressing change every 7 days and as needed; change immediately if damp, loosened, or visible soiled Perform Lab draw per: Lab orders: (Select all that apply)											
LABS	Perform weekly lab	Lab draw per: (Select one)		•		DI INI	- CDI		CDD	- FCD		
	draw on	☐ Home Health		□ BMF		BUN	□ CPK		□ CRP	□ ESR		
	Mondays,		□ CBC w/			Creatinine						
	as follows:					ripheral venipuncture, prior to dose then weekly						
	Fax lab results to:						of North Tex	as [□ Provider	s office		
*Product selection permitted unless dispense as written checked or clearly written on order												
										ENICE AC MIDIT	TTENI	
PROV	/IDER SIGNATUR	RE			D	ATE/TIME	 :		שוט וו	ENSE AS WRIT	IIEN	
					_	,						

Name:	Date of Birtl	n: Date:						
PICC Line placement for long-term antibiotics								
Dx:								
	Dispense as written	Substitution Permitted						



601 S. Main St., Suite 110, Keller, TX 76248 P / 682.357.4040 www.vitalcarenorthtexas.com